

sides and in waste ground, especially in those parts of the country which have been longest settled. It is usually found in bunches, and attains the full height of a man. It flowers from June until the arrival of frost.

The very natural order, called Umbellatæ by Linnæus and Umbelliferae by Jussieu, to which this plant and the following one belong, is found in the class Pentandria and order Digynia of the Linnæan artificial method.

The genus Conium of Linnæus has *both general and partial involucre*, the latter being *halved*. The fruit roundish and furrowed.

The species maculatum has the fruit *unarmed with the ridges undulated*.

Its more complete description is as follows. Root biennial, somewhat fusiform and generally branched. Stalk round, very smooth, striated, hollow, jointed, and more or less marked with purplish spots. Leaves two or three times pinnate, of a very bright green, with long, sheathing petioles inserted on the joints of the stem; the leaflets pinnatifid and toothed. Flowers in terminal umbels, the general involucre with half a dozen lanceolate, reflected leaflets, the partial involucre with three or four situated on the outside. Flowers very small, white. Petals five, oval with their

points inflexed. Stamens five, spreading, about the length of the corolla. Germ inferior. Styles two reflexed outwardly. Fruit roundish-oval, compressed, ribbed, the ribs being transversely wrinkled or crenate; separating into two oblong-hemispherical seeds.

Hemlock when fresh has a strong nauseous odour and taste. If the green leaves are distilled, the water which collects in the receiver has an insupportably nauseous taste, while that which remains in the retort is comparatively insipid. This circumstance, and likewise the fact that the dried leaves become inert by age and exposure, render it probable that the chief medicinal efficacy resides in a volatile portion of the plant. Of the more fixed ingredients of Hemlock, a variety of analyses have been made. The most recent which I have met with is that of Schrader, who from a thousand grains of the plant obtained the following substances. Extractive 27.3—Gummy extract 35.2—Resin 1.5—Albumen 3.4—Green fæcula 8.—He also detected various earthy and alkaline salts. These however are found to vary according to the soil in which the plant grows. The volatile portion, which I obtained in water distilled from the leaves, did not exhibit any essential oil, and effected no change in the colour of litmus. It